

Medical Standing Order for NASAL NALOXONE 4 mg

2022 - 2023 School Year

Opioid Overdose - Always an EMERGENCY

- Naloxone (Narcan) is indicated for reversal of suspected opioid overdose resulting in respiratory depression and/or unresponsiveness to stimuli.
- Person with symptoms of overdose must ALWAYS be transported by EMS to hospital for additional medical attention.

Signs of Overdose

- Neuro Unresponsive to stimuli such as shaking or sternal rub, limp body.
 Pinpoint Pupils
- <u>Lungs</u> slow, shallow breathing or no breathing.
- <u>Heart</u> Slow or no pulse. Cyanotic coloration to lips and fingertips.
- Mouth unable to speak/incoherent. Gurgling or deep snorting.
- Skin Pale, Clammy. Cyanotic coloration to lips and fingertips.
- **GI** Vomiting or gurgling.
- Other Report from bystanders, prior history or presence of drug paraphernalia

Action:

- Call 911
- Institute School Emergency Plan
- Administer NARCAN Nasal Spray
 - > Lay person on back
 - > Tilt head back slightly
 - > Insert Narcan nozzle tip gently into ONE NOSTRIL
 - > Press Plunger Firm to administer dose
 - > Place person in Recovery position



- Repeat Dose in 3 5 minutes to other nostril if person does not respond by waking up to voice or touch.
- Monitor vital signs and airway until EMS arrive; if needed start Rescue breathing and CPR
- Individual is to be transported by EMS even if symptoms improve.
- Continue to monitor vital signs and record. Provide to EMS along with a copy of student's Emergency Form / permission to treat.
- If the parents/guardian are not present at time of transport to the hospital, a school representative must accompany the student to the hospital.

Mil Stein, mos		4/1/2022	
Neil Stein, MD	Medical Director	Date	